

30753 to 30763—Continued.**30760. GLORIOSA SUPERBA L.**

Distribution.—Throughout tropical India from the northwestern Himalayas to Ceylon and eastward to Malakka and Cochin China; also in the Mozambique district of eastern Africa and in Guinea on the western coast.

30761. CAJUPUTI HYPERICIFOLIA (Salisb.) Skeels.

(*Metrosideros hypericifolia* Salisb. Prodrumus, 1796, p. 351).

The seeds of this Australian shrub belonging to the family Myrtaceæ were received under the name *Melaleuca hypericifolia* Smith (Transactions of the Linnean Society, vol. 3, 1797, p. 279). The generic name *Melaleuca* was published by Linnæus (Mantissa Plantarum, p. 105) in 1767, with one species, *M. leucadendra*, based on *Myrtus leucadendra* Stickman (Herbarium Amboinense, 1754, p. 9; reprinted in Linnæus's *Amoenitates Academicæ*, vol. 4, 1759, p. 120). However, in 1763, Adanson (*Familles des Plantes*, vol. 2, pp. 84, 530) had published the generic name *Cajuputi*, based on plate 16 of volume 2 in Rumph's *Herbarium Amboinense*, the same plate on which *Myrtus leucadendra* was founded. It is therefore necessary to use the generic name *Cajuputi* for the species congeneric with *Myrtus leucadendra* Stickman. This course has already been adopted by Lyons (*Plant Names, Scientific and Popular*, 1900, p. 74); see No. 30793.

Distribution.—A tall shrub with rich red flowers in spikes 2 inches long, found in damp places in the vicinity of Port Jackson in New South Wales, Australia.

30762. PRIMULA BULLEYANA Forrest.**Primrose.****30763. PRIMULA LITTONIANA Forrest.****Primrose.****30764 to 30822.**

From Cambridge, England. Presented by Dr. R. Irwin Lynch, curator, University Botanic Garden, Cambridge University. Received April 25, 1911.

Seeds of the following:

30764. CLEMATIS INTEGRIFOLIA × VITICELLA.**Clematis.**

See No. 29927 for previous introduction.

30765. CLEMATIS GRAVEOLENS Lindl.**Clematis.**

Distribution.—On the temperate slopes of the western Himalayas at an elevation of 6,000 to 11,000 feet in northern India.

30766. CLEMATIS HERACLEAEFOLIA DC.**Clematis.**

Distribution.—On the slopes of the mountains in the provinces of Chihli and Shantung, China.

30767. CLEMATIS INTEGRIFOLIA L.**Clematis.**

See No. 28844 for previous introduction.

30768. CLEMATIS RECTA L.**Clematis.**

Distribution.—Southern Europe extending from northern Spain to central Russia.

30769. BERBERIS LYCIUM Royle.**Barberry.**

See No. 1603 for previous introduction.

Distribution.—On the dry, hot slopes of the Himalayas at an elevation of 3,000 to 9,000 feet in the northwestern part of India.

30770. × BERBERIS STENOPHYLLA Lindl.