

32175 to 32245—Continued.**32194—Continued.**

is generally called *Entada scandens*. This use of the generic name *Lens* by Stickman prevents the use of the same name for the lentil. As no other generic name had been published for the lentil up to 1909, Mr. W. F. Wight (Century Dictionary Supplement, p. 719) published the name *Lentilla* by referring to "lens" and "lentil," under which names on pages 3409 and 3410 of volume 4 of the 1895 edition the lentil is described as *Lens esculenta*. When the new edition of 1911 appeared, in which the information first published in the two supplementary volumes 11 and 12 of 1909 was incorporated with the main body of the work, the name *Lentilla lens* was published (vol. 5, p. 3410) under the heading "lentil."

From Barnaul, Siberia.

"(No. 1652a, August 31, 1911.) A small-seeded variety of lentil that is grown by some progressive farmers in southern Siberia as a fodder for horses and milch cows. Prefers a light, sandy soil. Of value possibly for forage purposes in the cooler sections of the semiarid belt." (*Meyer.*)

32195. VICIA SATIVA L.**Spring vetch.**

From Barnaul, Siberia.

"(No. 1653a, August 31, 1911.) A vetch that is grown here and there in Siberia for forage purposes. It is said to stand drought very well and promises to be a good fodder plant in semiarid southwestern Siberia." (*Meyer.*)

32196. VICIA CRACCA L.**Vetch.**

From Tomsk, Siberia.

"(No. 1654a, August 18 to 20, 1911.) A wild vetch found on slightly shaded places. Likes to overrun low scrub. Is eagerly eaten by horses and cattle. Of value in pastures of the northern United States where some scrub is scattered over the land." (*Meyer.*)

32197. VICIA AMOENA Fisch.**Vetch.**

From near Chistunka, southwestern Siberia.

"(No. 1655a, September 8 and 9, 1911.) A sturdy vetch with rather dark green foliage found on dry earth banks. Does not grow over 1 foot in height, but on favorable locations spreads out considerably. Of value possibly as a forage or pasture plant in the northern sections of the United States." (*Meyer.*)

32198. VICIA AMOENA Fisch.**Vetch.**

From near Barnaul, Siberia.

"(No. 1656a, September 1 to 3, 1911.) A wild vetch, occurring along the edges of abandoned fields. Apparently a variety of the preceding number and as such the same remarks apply to it." (*Meyer.*)

32199. VICIA MEGALOTROPIS Ledeb.**Vetch.**

From near Sminogorsk, southwestern Siberia.

"(No. 1657a, September 21 to 24, 1911.) An exceedingly vigorous species of vetch, often 8 to 10 feet in height when running up into bushes; in open places, however, it reaches only a couple of feet in height. Much relished by horses and cattle and apparently of great promise as a forage plant. To be tested in the cooler sections of the United States." (*Meyer.*)

Distribution.—Slopes of the mountains in the Altai region of southwestern Siberia.